

Ram pressure stripping of halo gas in disk galaxies: Implications on galactic star formation in different environments

Kenji Bekki^{1*}

¹*School of Physics, University of New South Wales, Sydney NSW, 2052, Australia*

Accepted, Received 2005 February 20; in original form

ABSTRACT

We numerically investigate evolution of gaseous halos around disk galaxies in different environments ranging from small groups to rich clusters in order to understand galaxy evolution in these environments. Our simulations self-consistently incorporate effects of ram pressure of intergalactic medium (IGM) on disk and halo gas of galaxies and hydrodynamical interaction between disk and halo gas so that mass fractions of halos gas stripped by ram pressure of IGM (F_{strip}) can be better estimated. We mainly investigate how F_{strip} depends on total masses of their host environments (M_{host}), galactic masses (M_{gal}), densities and temperature of IGM (T_{IGM} and ρ_{IGM} , respectively), relative velocities between IGM and galaxies (V_r), and physical properties of disks (e.g., gas mass fraction). We find that typically 60–80% of halo gas can be efficiently stripped from Milky Way-type disk galaxies by ram pressure in clusters with $M_{\text{host}} \sim 10^{14} M_{\odot}$. We also find that F_{strip} depends on M_{host} such that F_{strip} is higher for larger M_{host} . Furthermore it is found that F_{strip} can be higher in disk galaxies with smaller M_{gal} for a given environment. Our simulations demonstrate that the presence of disk gas can suppress ram pressure stripping of halo gas owing to hydrodynamical interaction between halo and disk gas. Ram pressure stripping of halo gas is found to be efficient (i.e., $F_{\text{strip}} > 0.5$) even in small and/or compact groups, if $\rho_{\text{IGM}} \sim 10^5 M_{\odot} \text{ kpc}^{-3}$ and $V_r \sim 400 \text{ km s}^{-1}$. Based on the derived radial distributions of remaining halo gas after ram pressure stripping, we propose that truncation of star formation after halo gas stripping can occur outside-in in disk galaxies. We suggest that although gradual truncation of star formation in disk galaxies can occur in groups, it can proceed less rapidly in comparison with cluster environments. We also suggest low-mass galaxies are likely to truncate their star formation more rapidly owing to more efficient halo gas stripping in groups and clusters.

Key words: galaxies:halo – galaxies:structure – galaxies:kinematics and dynamics

1 INTRODUCTION

Since Larson et al. (1980) discussed removal of galactic halo gas from disk galaxies in terms of transformation from spirals into S0s, many observational and theoretical works investigated evolution of galactic halo gas in variously different aspects of galaxy evolution, such as maintenance of spiral arms by halo gas infall (e.g., Sellwood & Carlberg 1984), formation of passive spiral galaxies in groups and clusters (e.g., Bekki et al. 2002), and color evolution of satellite galaxies entering into their host group- and cluster-scale halos (e.g., Font et al. 2008). One of important suggestions from

these previous works is that stripping of galactic halo gas can cause severe suppression of galactic global star formation in different environments (“strangulation”; Balogh et al. 1999, 2000). Although the strangulation scenario can explain a number of recent observational results of galaxies, such as the presence of red passive spirals in distant clusters (e.g., Couch et al. 1998, Dressler et al. 1999, Poggianti et al. 1999, 2008) and mean star formation rate dependent on galaxy environments (e.g. Balogh et al. 2004), it is still less clear why and how effective strangulation can occur in different environments (e.g., groups and clusters).

Previous theoretical and numerical studies tried to understand how galactic halo gas responds to environmental effects, such as tidal fields of groups and clusters (e.g., Bekki

* E-mail: bekki@phys.unsw.edu.au

et al. 2001) and ram pressure stripping of IGM (e.g., Balogh et al. 2000; Bekki et al. 2002; Hester 2006; McCarthy et al. 2008). Although these works discussed quantitatively how much fraction of halo gas can be removed from galaxy-scale halos by ram pressure of IGM in their host environments, their models are not so sophisticated at some points. For example, previous numerical models (e.g., Bekki et al. 2002) do not adopt realistic galaxy models with stellar and gaseous disks and ignores hydrodynamical interaction between halo and disk gas. More sophisticated numerical models are required to discuss more qualitatively how much fraction of halo gas can be removed by ram pressure stripping and whether and how hydrodynamical interaction between halo and disk gas can influence ram pressure stripping of halo gas in disk galaxies.

The purpose of this paper is thus to investigate time evolution of galactic halo gas in disk galaxies under moderately strong ram pressure of IGM in their host environments based on more sophisticated numerical simulations. We focus mainly on (i) final mass fractions of halo gas stripped from disk galaxies (F_{strip}), (ii) radial properties of halo gas, and (iii) dependences of (i) and (ii) on physical properties of galaxy environments (e.g., total masses of groups and clusters). The present results can be used for interpreting observational results on spatial distribution of hot halo gas in galaxies (e.g., Jeltama et al. 2008), physical properties of passive spirals in groups and clusters (e.g., Poggianti et al. 2008; Bamford et al. 2009), and origin of possible suppression of star formation in compact groups (e.g., Rasmussen et al. 2008).

The plan of the paper is as follows: In the next section, we describe our numerical models for ram pressure stripping of galactic halo gas. In §3, we present the numerical results mainly on the physical properties of remaining halo gas after the ram pressure stripping for variously different models. In §4, we compare the present results with those from other authors and discuss implications of the present results. We summarize our conclusions in §5. We here do not discuss ram pressure stripping of *disk gas* in galaxies in an extensive manner, because a number of authors have already discussed the stripping processes in detail (e.g., Abadi et al. 1999; Kronberger et al. 2008; Marcoline et al. 2003; Roediger & Bruggen 2008; Tonnesen & Bryan 2008; Vollmer et al. 2006).

2 THE MODEL

2.1 Disk galaxy

In order to simulate the time evolution of galactic halo gas under ram pressure of IGM, we use the latest version of GRAPE (GRAVity PipE, GRAPE-7) which is the special-purpose computer for gravitational dynamics (Sugimoto et al. 1990). We use our original GRAPE-SPH code (Bekki & Chiba 2006) which combines the method of smoothed particle hydrodynamics (SPH) with GRAPE for calculations of three-dimensional self-gravitating fluids in astrophysics. The original code used in our previous studies is here revised so that both (i) ram pressure effects of hot IGM on galactic halo gas and (ii) hydrodynamical interaction between halo and disk gas can be investigated in a fully self-consistent manner. Fig. 1 illustrates initial configurations for halo and




Figure 1. A schematic representation of the present numerical model for ram pressure stripping of galactic halo gas in disk galaxies. Both spherical gaseous halo and initially thin gaseous disk in a disk galaxy can be influenced by hot IGM in groups and clusters of galaxies in the present model. Both hydrodynamical interaction between halo and disk gas and that between hot IGM and disk and halo gas are modeled in a self-consistent manner in the present study.

disk gas in a disk galaxy and hot IGM surrounding the disk galaxy in the present numerical study.

Since our numerical methods for modeling dynamical evolution of Milky Way-type disk galaxies have already been described by Bekki & Shioya (1998) and by Bekki & Peng (2006), we give only a brief review here. The total disk mass and the size of a disk of a Milky Way-type disk galaxy with the total mass of M_{gal} are M_d and R_d , respectively. Henceforth, all masses and lengths are measured in units of M_d and R_d , respectively, unless specified. Velocity and time are measured in units of $v = (GM_d/R_d)^{1/2}$ and $t_{\text{dyn}} = (R_d^3/GM_d)^{1/2}$, respectively, where G is the gravitational constant and assumed to be 1.0 in the present study. If we adopt $M_d = 6.0 \times 10^{10} M_\odot$ and $R_d = 17.5$ kpc as a fiducial value, then $v = 1.21 \times 10^2$ km/s and $t_{\text{dyn}} = 1.41 \times 10^8$ yr, respectively. The disk is composed of a dark matter halo, a stellar disk, a stellar bulge, a gaseous disk, and a gaseous halo.

The mass ratio of the dark matter halo to the stellar disk in a disk model is fixed at 16.7 models (i.e., $M_{\text{gal}}/M_d = 17.7$). We adopt the density distribution of the NFW halo (Navarro, Frenk & White 1996) suggested from CDM simulations:

$$\rho(r) = \frac{\rho_0}{(r/r_s)(1+r/r_s)^2}, \quad (1)$$

where r , ρ_0 , and r_s are the spherical radius, the characteristic density of a dark halo, and the scale length of the halo, respectively. The value of r_s (0.6 in our units for $c = 10$) is chosen such that the rotation curve of a disk is reasonably consistent with observations. The mass fraction and the scale length of the stellar bulge represented by the Hernquist profile are fixed at 0.17 (i.e., 17 % of the stellar disk) and 0.04 (i.e., 20 %) of the scale length of the stellar

Table 1. The ranges of model parameters.

Parameters	$M_{\text{gal}} (\times M_{\text{MW}})^a$	$V_r (\text{km s}^{-1})^b$	$\rho_{\text{IGM}} (\times 10^5 M_{\odot} \text{ kpc}^{-3})^c$	$T_{\text{IGM}} (\times 10^7 \text{ K})^d$	f_g^e	$R_{\text{halo}} (\times R_d)^f$
Value ranges	0.05–1.0	160–2000	0.1–5.0	0.1–3.2	0–0.1	1–6

^a The total mass of a galaxy in units of $M_{\text{MW}} (= 1.1 \times 10^{12} M_{\odot})$.

^b The relative velocity of the hot IGM with respect to the galaxy.

^c The initial mass density of the hot IGM in the cube.

^d The initial temperature of the hot IGM in the cube.

^e The initial gas mass fraction in a galactic disk.

^f The initial radius of a spherical galactic halo gas of a disk galaxy in units of R_d , where R_d is the disk size.




Figure 2. Morphological evolution of halo (magenta) and disk (cyan) gas projected onto the x - z plan in a disk galaxy for the standard model. The dark matter halo, stellar disk, bulge, and hot IGM are not shown for clarity. The time T in units of Gyr is shown in the upper left corner for each frame.

disk), respectively, which are consistent with those of the bulge model of the Galaxy.

The radial (R) and vertical (Z) density profiles of the disk are assumed to be proportional to $\exp(-R/R_0)$ with scale length $R_0 = 0.2$ and to $\text{sech}^2(Z/Z_0)$ with scale length $Z_0 = 0.04$ in our units, respectively: both stellar and gaseous disks follow this exponential distribution. In addition to the rotational velocity caused by the gravitational field of disk, bulge, and dark halo components, the initial radial and azimuthal velocity dispersions are assigned to the disc component according to the epicyclic theory with Toomre's parameter $Q = 1.5$. The vertical velocity dispersion at given radius is set to be 0.5 times as large as the radial velocity dispersion at that point, as is consistent with the observed trend of the Milky Way (e.g., Wielen 1977).

We investigate models with different M_{gal} and adopt the Freeman's law (Freeman 1970) to determine R_0 of a disk galaxy according to its disk mass:

$$R_0 = 3.5 \left(\frac{M_d}{6 \times 10^{10} M_{\odot}} \right)^{0.5} \text{ kpc}. \quad (2)$$

Structural and kinematical properties of dark matter halos and stellar disks are assumed to be self-similar between models with different M_{gal} . The gas mass fraction (f_g) is assumed to be a free parameter. An isothermal equation of state is used for the gas with temperatures of 10^4 K for models with $M_d = 6 \times 10^{10} M_{\odot}$. The initial temperature (T_{iso}) of disk gas is assumed to be scaled to $T_{\text{iso}} \propto M_{\text{gal}}^{0.5}$.

2.2 Halo gas and hot IGM

The gaseous halo has mass M_{hg} and the same spatial distribution as the dark matter and is assumed to be initially in hydrostatic equilibrium. The initial gaseous temperature of a halo gas particle is therefore determined by the gas density, total mass, and gravitational potential at the location of the particle via Euler's equation for hydrostatic equilibrium (e.g., the equation 1E-8 in Binney & Tremaine 1987). Therefore gaseous temperature $T_{\text{halo}}(r)$ at radius r from the center of a disk galaxy can be described as:

$$T_{\text{halo}}(r) = \frac{m_p}{k_B \rho_{\text{halo}}} \int_r^\infty \rho_{\text{halo}}(r) \frac{GM(r)}{r^2} dr, \quad (3)$$

where m_p , G , and k_B are the proton mass, the gravitational constant, and the Boltzmann constant, respectively, and $M(r)$ is the total mass within r determined by the adopted mass distributions of dark matter and baryonic components in the disk galaxy. Radiative cooling is not included in the present study so that the hydrodynamical equilibrium of halo gas can be obtained in isolated disk models: if gaseous cooling is included in a disk model, the halo gas can rapidly collapse to settle down onto the gas disk so that physical roles of ram pressure stripping in evolution of halo gas can not properly investigated. The present idealized models thus help us to grasp some essential ingredients of the physical roles of hot IGM in evolution of galactic halo gas.

Sembach et al. (2003) showed that the Galaxy gaseous halo is highly extended ($R \sim 70$ kpc) and low-density ($\rho_{\text{halo}} \approx 10^{-4} - 10^{-5} \text{ cm}^{-3}$). It is however observationally unclear how far galactic halo gas extends in other galaxies. We therefore consider that the radius of the galactic halo (R_{halo}) is a free parameter. Guided by the above observational results by Sembach et al. (2003), we mainly investigate models with $R_{\text{halo}} = 3 - 6R_d$ and $\rho_{\text{halo}} = 10^{-4} \text{ cm}^{-3}$.

The disk galaxy is assumed to be embedded in hot IGM with the temperature of T_{IGM} and the densities of ρ_{IGM} . In order to avoid huge particle numbers to represent the entire IGM in group and clusters of galaxies (e.g., Abadi et al. 1999), the IGM is represented by SPH particles with velocities of V_r in a cube with the size of R_{IGM} . The initial velocity of each SPH particle for the IGM is set to be $(V_r, 0, 0)$ for all models (i.e., the particle flows along the x -axis to the positive x direction). The IGM has a uniform distribution within the cube and R_{IGM} is set to be $10M_d$ for models with $R_{\text{halo}} = 3R_d$. We include periodic boundary conditions (at R_{IGM}) for the IGM SPH particles leaving the cube. The parameter values of T_{IGM} and ρ_{IGM} are chosen based on the adopted total masses of galaxy's host environment (M_{host}).

The spin of the disk galaxy is specified by two angles θ and ϕ (in units of degrees), where θ is the angle between the z -axis and the vector of the angular momentum of the disk, and ϕ is the azimuthal angle measured from x axis to the projection of the angular momentum vector of the disk onto the x - y plane. In order to show more clearly the ram pressure effects on galactic halo gas, we mainly investigate the models with $\theta = 90^\circ$ and $\phi = 0^\circ$, in which the entire gas disk can strongly feel ram pressure of IGM. We also investigate the models with different θ and ϕ to clarify the roles of halo-disk hydrodynamical interaction in keeping halo gas within disk galaxies.

The mass resolution for disk gas particles, halo ones, and IGM in luminous disk models with $M_d = 6 \times 10^{10} M_\odot$ are $3.5 \times 10^5 M_\odot$, $5.9 \times 10^4 M_\odot$ and $5.3 \times 10^5 M_\odot$, respectively. The typical smoothing length for disk gas particles (the mean smoothing length at $T = 0.21$ Gyr) in the standard model are 680 pc (this can be as large as 4.7 kpc for halo gas particles owing to the stripped halo gas).

2.3 Parameter study

Although we have investigated 45 models with different M_{gal} , f_g , V_r , ρ_{IGM} , and T_{IGM} , we mainly show the results of

the “standard” model with $M_{\text{gal}} = 1.1 \times 10^{12} M_\odot$, $f_g = 0.1$, $V_r = 500 \text{ km s}^{-1}$, $\rho_{\text{IGM}} = 10^5 M_\odot \text{ kpc}^{-3}$, and $T_{\text{IGM}} = 10^7 \text{ K}$. This is mainly because the standard model can clearly show a typical behavior of ram pressure stripping of halo gas from disk galaxies. For a canonical baryonic mass fraction of 0.14 ($= \Omega_b/\Omega_m$) in the universe, ρ_{IGM} can be as high as $10^5 M_\odot \text{ kpc}^{-3}$ at $r \sim 100$ kpc in a cluster with the NFW profile, the total mass of $10^{14} M_\odot$, and the virial velocity (v_{vir}) of $500 - 600 \text{ km s}^{-1}$. We thus consider that the adopted values of ρ_{IGM} , V_r , and T_{IGM} are reasonable for a cluster of galaxies with $M_{\text{host}} = 10^{14} M_\odot$. It should be stressed here that the strength of ram pressure (P_{ram}) in the standard model is much weaker than that required for ram pressure stripping of disk gas in the *central regions of clusters* (e.g., Abadi et al. 1999). The adopted galaxy mass in the standard model is referred to as M_{MW} just for convenience.

The total number of particles used for the disk galaxy and the IGM in the standard model are 133699 and 125000, respectively. The present simulation requires typically 70 – 160 CPU hours of the adopted GRAPE-7 systems for each model depending on model parameters: we have adopted the above particle numbers to run numerous models within a reasonable time scale. We have conducted a resolution test by using a high-resolution model with the total particle number of 333805 and found that there is only $\sim 7\%$ difference in F_{strip} between the standard model and the high-resolution one. We thus consider that the above particle number is enough to discuss ram pressure stripping of halo gas around disk galaxies. Gravitational softening lengths in simulation units (R_d) are set to be fixed at 0.018 for the galaxy and 0.2 for the IGM. The range of model parameters investigated in the present study are shown in the Table 1.

We divide models into four categories: “Rich cluster” model with $M_{\text{host}} = 10^{15} M_\odot$, “cluster” one with $M_{\text{host}} = 10^{14} M_\odot$, “group” one with $M_{\text{host}} = 10^{13} M_\odot$, and “small group” one with $M_{\text{host}} = 10^{12} M_\odot$. Considering virial velocities (v_{vir}) of dark matter halos (NFW95) and the scaling relation between masses and sizes for dark matter halos (e.g., Padmanabhan 1993), reasonable values of T_{IGM} and V_r are chosen for the above four models with different M_{host} . In the present study, V_r of IGM in a galaxy environment with M_{host} is set to be similar to v_{vir} of a dark matter halo with M_{host} . T_{IGM} in the small group, group, cluster, and rich-cluster models are 10^6 K , $3.2 \times 10^6 \text{ K}$, 10^7 K , and $3.2 \times 10^7 \text{ K}$, respectively, in the present study. V_r in the small group, group, cluster, and rich-cluster are 158 km s^{-1} , 281 km s^{-1} , 500 km s^{-1} , and 889 km s^{-1} , respectively.

We mainly investigate the time evolution of gas mass (M_g) for halo and disk components in galaxies and the final mass fraction of halo gas stripped from galaxies (F_{strip}). We consider that if SPH gas particles at the final time step of a simulation are outside R_{halo} , they are regarded as being stripped by ram pressure of IGM. For most models, halo gas can be rapidly stripped from galaxies within well less than 0.5 Gyr. Therefore, we estimate F_{strip} at $T = 0.56$ Gyr, where the time T represents the time that has elapsed since the simulation starts. We also estimate the accretion rate of gas onto the central 1 kpc of a disk galaxy in a simulation by dividing the total gas mass accumulated in the central 1 kpc at the final time step by time that has elapsed since the simulation starts (i.e., 0.56 Gyr). We confirm that the




Figure 3. Time evolution of the total mass ($M_g(T)$) of disk (solid) and halo (dotted) gas normalized to their initial values ($M_g(T=0)$) in the standard model.




Figure 4. The initial distances ($R_{\text{hg,ini}}$) of halo SPH particles as a function of the final ones ($R_{\text{hg,fin}}$) at $T = 0.56$ Gyr (i.e., final time step) for the standard models. A big filled circle represents the mean value of $R_{\text{hg,ini}}$ for each $R_{\text{hg,fin}}$ bin. The dotted line shows $R_{\text{hg,fin}} = R_{\text{hg,ini}}$. Since SPH particles that can finally remain within the galaxy ($R_{\text{hg,fin}} < 3R_d$) are plotted in this figure, the result means that the vast majority of the remaining halo gas after ram pressure stripping can originate from the inner halo ($R_{\text{hg,ini}} \sim 10$ kpc). This suggests that truncation of star formation after ram pressure stripping of halo gas can proceed outside-in.

accretion rates in different models do not depend so much on model parameters in the present study.

3 RESULTS

3.1 The standard model

Fig. 2 shows how halo gas responds to moderately strong ram pressure of IGM in the standard model with $M_{\text{host}} = 10^{14} M_{\odot}$, $T_{\text{IGM}} = 10^7$ K, and $V_r = 500$ km s $^{-1}$. The halo gas is efficiently stripped from the disk galaxy to start to form a gaseous stream behind the galaxy within 0.14 Gyr. Owing to the presence of the disk gas, halo gas initially located in




Figure 5. The cumulative mass distribution of halo (red) and disk (blue) gas at $T = 0$ Gyr (solid) and $T = 0.56$ Gyr (dotted) in the standard model. The masses at R are normalized to those at $R = 3R_d$.




Figure 6. The same as Fig.3 but for the group model.

the inner halo can be accumulated above the disk and thus can not be stripped from the galaxy. The gaseous stream with many small clumps formed after stripping can finally become less remarkable within 0.56 Gyr and most of the remaining halo gas can be located close to the disk. The final distribution of the halo gas appears to be more compact, flattened, and inhomogeneous at $T = 0.56$ Gyr. The disk gas, on the other hand, can not be stripped efficiently from the galaxy, because it is much more strongly bounded by the




Figure 7. Final distributions of halo (magenta) and disk (cyan) gas projected onto the x - z plane for the group model with $M_{\text{host}} = 10^{13} M_{\odot}$ (left) and the cluster one with $M_{\text{host}} = 10^{14} M_{\odot}$ (right). The green solid circle in each frame represents the initial size of the gaseous halo.

disk in comparison with the halo gas. Thus, ram pressure stripping of halo gas is much more effective than that of disk gas.

Fig. 3 shows that 66% and 6% of initial gas can be stripped from halo and disk, respectively, by ram pressure within 0.56 Gyr for the standard model. Fig. 4 shows that halo gas initially in the inner halo ($R_{\text{hg,ini}} < 10$ kpc) is more likely to be still located in the galaxy ($R_{\text{hg,fin}} < R_{\text{halo}}$). These remaining halo gas might well be accreted later onto the inner region of the disk to increase slightly the total gas mass of the disk. The halo gas initially located in the outer halo ($R > R_d$) can be much more efficiently stripped by ram pressure. The stripping of the outer halo gas might well severely suppress the accretion of the gas onto the outer part of the galaxy and thus contribute to truncation of star formation there.

Fig. 4 shows that the radial distribution of the halo gas can dramatically change after hydrodynamical interaction between IGM and the halo gas, though that of the disk gas can hardly change owing to ram pressure weaker than the self-gravity of the gas disk. The distribution of halo gas becomes more compact with the half-mass radius of ~ 10 kpc at $T = 0.56$ Gyr, because the stripping can happen much more efficiently in the outer halo, where the halo is more susceptible to ram pressure effect owing to weaker gravity of the galaxy there. These results in Figs. 2–5 clearly demonstrate that moderately strong ram pressure of IGM can very efficiently remove the halo gas, even though it can not remove the disk gas.

In order to understand more clearly how the disk gas in a disk galaxy can promote or suppress the ram pressure stripping of the halo one, we have investigated a comparative model with no disk gas (“no disk gas model”, $f_g = 0$). It is found that F_{strip} in the no disk gas model is 0.79, which is by a factor of 1.2 higher than that (0.66) in the standard model with disk gas ($f_g = 0.1$). The final distribution of the halo gas in the no disk gas model is less flattened and a thin gaseous stream behind the disk can not be seen. These results mean that hydrodynamical interaction between disk and halo gas can suppress ram pressure stripping of the halo gas.

As shown in Fig. 2, the remaining halo gas at $T = 0.56$ Gyr is compressed and accumulated above the gas disk. About 28% of the remaining halo gas can be located at $X < 0$ (i.e., left side of the gas disk) and is likely to be accreted finally onto the disk for further star formation: the low-density gas located at $X > 0$ (i.e., right side of the disk) is less likely to be accreted rapidly onto the disk. Therefore, the mass fraction of halo gas that can be used for later star formation (F_{sf}) after ram pressure stripping can be significantly smaller than $1 - F_{\text{strip}}$ in disk galaxies (F_{sf} can be typically $\sim 0.3 \times (1 - F_{\text{strip}})$).

Interestingly, about 0.7% of the halo gas (corresponding roughly to $1.1 \times 10^7 M_\odot$ within R_{halo}) can be fueled to the central 1kpc of the disk. This radial transfer of halo gas to the central region of the galaxy can be seen in other models with different model parameters. These implies that if the centrally accumulated halo gas can be further transferred to the vicinity of the massive black hole (MBH) located in the center of the bulge, the halo gas can be used for fueling the MBH with the accretion rate of $\sim 0.02 M_\odot \text{ yr}^{-1}$. These




Figure 8. Parameter dependences of F_{strip} on V_r (upper left), M_{host} (upper right), M_{gal} (lower left), and ρ_{IGM} (lower right). For each parameter dependence, model parameters other than the described one (e.g., V_r in the upper left frame) are set to be the same as those used in the standard model. In the lower right panel, $\rho_{\text{IGM},0}$ is the same as $\rho_{\text{IGM}} (= 10^5 M_\odot \text{ kpc}^{-3})$ used in the standard model.




Figure 9. The same as Fig. 3 but for four different models with different M_{gal} in the group model with $M_{\text{host}} = 10^{13} M_\odot$: $M_{\text{gal}} = 0.05 M_{\text{MW}}$ (solid), $M_{\text{gal}} = 0.1 M_{\text{MW}}$ (dotted), $M_{\text{gal}} = 0.5 M_{\text{MW}}$ (short-dashed), and $M_{\text{gal}} = 1.0 M_{\text{MW}}$ (long-dashed).

results imply that ram pressure of IGM can be responsible for the activation of weak active galactic nuclei (AGN).

3.2 Parameter dependences

The dependences of F_{strip} and final radial distributions of halo gas are described as follows:

(i) Galactic halo gas can be efficiently stripped from disk galaxies by ram pressure of IGM in groups with $M_{\text{host}} \approx 10^{13} M_\odot$, though F_{strip} can be smaller than those in cluster models with $M_{\text{host}} \approx 10^{14} M_\odot$. For example, as shown in Fig. 6, the group model with $M_{\text{host}} = 10^{13} M_\odot$ shows $F_{\text{strip}} = 0.38$, which is significantly smaller than that (0.66) in the cluster model with $M_{\text{host}} = 10^{14} M_\odot$. Fig. 7 clearly shows that the final distribution of halo gas in the group model is less compact and less flattened than that in the cluster model owing to weaker effects of ram pressure of IGM in the group model.

(ii) F_{strip} depends on M_{host} in such a way that F_{strip}




Figure 10. The same as Fig. 3 but for four different models with different V_r in the cluster model with $M_{\text{host}} = 10^{14} M_{\odot}$: $V_r = 500 \text{ km s}^{-1}$ (solid), $V_r = 1000 \text{ km s}^{-1}$ (dotted), $V_r = 1500 \text{ km s}^{-1}$ (short-dashed), and $V_r = 2000 \text{ km s}^{-1}$ (long-dashed).

is higher in models with higher M_{host} . As shown in Fig. 8, the models with $M_{\text{host}} = 10^{12} M_{\odot}$ (corresponding to small groups like the Local Group) do not show any efficient ram pressure stripping of halo gas. This result suggests that ram pressure stripping is not so important for galactic global star formation of luminous disk galaxies in small groups with masses of $\sim 10^{12} M_{\odot}$.

(iii) Less massive disk galaxies are likely to lose a larger amount of their halo gas more rapidly for a given environment, though the dependence of F_{strip} on M_{gal} is not very strong. (See Fig. 8). Fig. 9 clearly shows that even in the group environment with $M_{\text{host}} = 10^{13} M_{\odot}$, less massive galaxies can lose larger mass fractions of their halo gas more rapidly. These results imply that less luminous disk galaxies can truncate more rapidly their star formation after they enter into group and cluster environments.

(iv) For a given M_{host} , F_{strip} are higher in models with higher V_r owing to stronger force of ram pressure of IGM (See Fig. 8). Also Fig. 10 shows that galactic halo gas can be more rapidly stripped from disk galaxies by ram pressure in models with higher V_r . These results suggest that disk galaxies passing through the inner regions of groups and clusters can truncate their star formation more rapidly owing to more efficient ram pressure stripping of their halo gas.

(v) F_{strip} depends on ρ_{IGM} such that F_{strip} is higher for models with higher ρ_{IGM} (See Fig. 8), which is a natural result of $P_{\text{ram}} \propto \rho_{\text{IGM}} \times V_r^2$. It is found that if $\rho_{\text{IGM}} \sim 10^5 M_{\odot} \text{ kpc}^{-3}$ and $V_r \sim 400 \text{ km s}^{-1}$ in the small group model with $M_{\text{host}} = 10^{12} M_{\odot}$ and $T_{\text{IGM}} = 10^6 \text{ K}$, then F_{strip} can be as high as 0.65. This result suggests that truncation of star formation by ram pressure stripping of halo gas can be possible even in small and/or compact groups, if their IGM densities are as high as $10^5 M_{\odot} \text{ kpc}^{-3}$ and member galaxies can have moderately larger velocity dispersion ($\sim 400 \text{ km s}^{-1}$).

(vi) F_{strip} does not depend strongly on f_g , θ , ϕ , and R_{halo} . For example, F_{strip} can differ only by a factor of ~ 1.2 between models with $f_g = 0.01 - 0.1$. Also F_{strip} can differ by a factor of ~ 1.2 between models with different θ and ϕ . These results imply that initial distributions of halo and disk gas are less important than other parameters (e.g., V_r) in determining F_{strip} .

4 DISCUSSION

4.1 Comparison with previous works

Bekki et al (2002) first investigated how much amount of galactic halo gas can be stripped by ram pressure in groups and clusters of galaxies based on rather idealized numerical simulations. They found that (i) about 90% of galactic halo gas can be stripped by the combination of tidal and ram pressure stripping in clusters of galaxies, (ii) F_{strip} depends on orbits of galaxies in groups and clusters, and (iii) galactic halo gas can be effectively removed from galaxies in groups only if the orbits of the galaxies are rather eccentric. Although their results can be useful for better understanding the origin of passive spirals, their models are not sophisticated enough to discuss the dependences of F_{strip} on galaxy environments in a quantitative manner.

Recently McCarthy et al. (2008) have conducted a thorough parameter study on ram pressure stripping of hot gaseous halos in galaxies for groups and clusters of galaxies. Their numerical simulations have shown that typically 70% of the initial halo gas around galaxies can be stripped by ram pressure within 10 Gyr in groups and clusters. Although they did not include stellar disks and bulges in their simulations, the result of F_{strip} is broadly consistent with our simulations ($F_{\text{strip}} \sim 0.6 - 0.8$), in which cosmologically motivated initial conditions of IGM and halo gas are not adopted. The slight difference in F_{strip} between their and our simulations would be due simply to the fact that they did not include disk gas, which is demonstrated to suppress ram pressure stripping of galactic halo gas in the present study. McCarthy et al. (2008) also have found that F_{strip} is higher in galaxies with smaller masses for a cluster environment with $M_{\text{host}} = 10^{14} M_{\odot}$ (i.e., the models with higher mass ratios show higher F_{strip} in their Fig. 8). This is also consistent qualitatively with our results.

Kawata & Mulchaey (2008) have investigated how ram pressure of a groups with $M_{\text{host}} = 8 \times 10^{12} M_{\odot}$ influences gaseous components in disk galaxies with maximum circular velocities of $\sim 150 \text{ km s}^{-1}$ based on cosmological chemodynamical numerical simulations. They have found that ram pressure in the group can be more than enough to remove hot halo gas of the galaxies, though it can not remove their cold gas within disks. These results are consistent qualitatively with the present ones, though initial conditions of disk galaxies in groups are different between Kawata & Mulchaey (2008) and the present study. This consistency implies that ram pressure of IGM in groups with $M_{\text{host}} \sim 10^{13} M_{\odot}$ can play an important role in controlling gas accretion from halos onto galactic disks and thus determining star formation histories within them.

Although broadly consistent results between the present and previous results clearly demonstrate that evolution of hot halo gas around galaxies under ram pressure of IGM can be one of key determinants for galaxy evolution, it is not so unclear how global properties of galaxies, in particular, their morphological, structural, and kinematical properties, change after removal of their halo gas. The present and previous simulations do not enable us to investigate the details of dynamical properties of disk galaxies after ram pressure stripping in a fully self-consistent manner. Therefore it is still unclear whether the observed rapid evolution of S0 fraction in groups (e.g., Wilman et al. 2009) can result

from truncation of star formation in spirals owing to the removal of halo gas in group environments. Given that minor and unequal-mass merging can create S0s (e.g., Bekki 1998), it is important for our future studies to confirm whether the simulated physical properties of S0s formed from spirals through removal of their halo gas can reproduce the observed ones reasonably well.

4.2 Implications of the present results

4.2.1 Outside-in truncation of star formation

The present study has demonstrated that outer halo gas can be preferentially stripped from disk galaxies during hydrodynamical interaction between hot IGM and halo gas: the remaining gas around disks mostly originates from inner halos. This result implies that if halo gas of a galaxy has intrinsic angular momentum to be accreted onto the disk, the preferential removal of the outer gas might well result in the suppression of the growth of the outer disk of the galaxy: the inner disk would grow after ram pressure stripping, though the accretion rate of halo gas would be significantly reduced. Our previous work (Bekki et al. 2002) showed that after removal of halo gas, stellar velocity dispersions (thus the Q parameter) of a disk galaxy can significantly increase in its outer part owing to dynamical heating of the disk by spiral arms. The present result combined with our previous one therefore suggests that gradual truncation of star formation can proceed outside-in in disk galaxies owing to more significantly increased Q and reduced f_g (i.e., gas mass fraction) in their outer disks.

This possible outside-in truncation scenario of star formation can provide a clue to the origin of dusty star-forming regions observed in passive spiral galaxies in distant groups and clusters (e.g., Dressler et al. 2009). Recent Spitzer observations of passive spiral galaxies have shown that passive spirals, which were previously suggested to be “dead and red” disk galaxies (Couch et al. 1998), show some levels of star formation (Dressler et al. 2009). The observed dusty star-forming, optically red spirals mean that star formation activities can be heavily obscured by dust. It is not clear why passive spirals, which should be gas-poor galaxies to explain the observed red colors (i.e., low-level star formation), can show heavily observed star-forming regions (which requires high-density gas and dust). The present outside-in truncation implies that star formation can continue only in the inner high-density regions (after stripping of halo gas), where dust extinction is highly likely to be significant. Thus the outside-in truncation scenario can naturally explain why passive spirals can have dusty-star forming regions without showing star-forming regions not obscured by dust in their outer parts.

Typically $\sim 5\%$ of halo gas can be accumulated above gas disks during ram pressure stripping and these gas can be strongly compressed and thus strongly interact hydrodynamically with the gas disks (see the Appendix A for the details for the evolution of gas disks during ram pressure stripping). As shown in previous numerical simulations (e.g., Bekki & Couch 2003; Kronberger et al. 2008), external high-pressure of gas may well trigger efficient star formation in gas disks. Therefore, the present results imply that before gradual truncation of star formation due to ram pressure

stripping of halo gas in disk galaxies, star formation in the galaxies could increase to some extent. Our future more sophisticated simulations including star formation within giant molecular clouds in disk galaxies under ram pressure of IGM will enable us to discuss this problem in a more quantitative way.

4.2.2 Recycling of ISM

Recent numerical simulations have shown that metal-rich gas ejected from massive OB stars and supernovae formed during efficient star formation in disk galaxies can be accreted onto the thin disks owing to hydrodynamical interaction between the gaseous ejecta and the gaseous halos (Bekki et al. 2009). This physical mechanism of galactic halo gas to keep gaseous ejecta within galaxies can effectively work only if the densities of gaseous halos can be as high as 10^{-5} cm^{-3} (Bekki et al. 2009). These results combined with the present ones therefore suggest that if galaxies enter inter group and cluster environments and lose most of their halo gas owing to ram pressure stripping of the halo gas, then recycling processes of metal-rich gaseous ejecta in interstellar medium (ISM) of galaxies can be dramatically changed owing to gaseous halos with much lower densities.

If recycling processes of metal-enriched ISM can be severely suppressed by ram pressure stripping of galactic halo gas, then chemical evolution of disk galaxies after the stripping can be also significantly changed. For example, energetic stellar winds from supernovae and massive OB stars in luminous disk galaxies under the influence of ram pressure can easily escape from the galaxies and be dispersed into IGM of their host environments. Less energetic winds like those from AGB stars, on the other hand, can be still trapped within the galaxies so that their ejecta can be used for further star formation and thus for chemical evolution. As a result of these, abundance patterns might well be significantly different between field disk galaxies, which can retain gaseous ejecta both from AGB stars and supernovae well, and cluster/group disk galaxies, which can retain only AGB ejecta. Our future quantitative investigation on chemical evolution of disk galaxies after ram pressure stripping of their halo gas will enable us to address this important question as to how abundance patterns in disks are different between field and group/cluster disk galaxies.

5 CONCLUSIONS

We have investigated time evolution of galactic halo gas in disk galaxies under moderately strong ram pressure of IGM in different environments based on self-consistent hydrodynamical simulations with variously different model parameters. We summarize our principle results as follows.

(1) Even moderately strong ram pressure of IGM with $\rho_{IGM} \approx 10^5 \text{ M}_\odot \text{ kpc}^{-3}$ and $V_r \approx 500 \text{ km s}^{-1}$ can strip galactic halo gas efficiently in clusters of galaxies with $M_{\text{host}} = 10^{14} \text{ M}_\odot$. Typically 60 – 80% of initial halo gas can be stripped from disk galaxies (i.e., $F_{\text{strip}} = 0.6 - 0.8$) in clusters, which is broadly consistent with results of other studies.

(2) F_{strip} depends on M_{host} such that it can be higher in environments with higher M_{host} . For example, F_{strip} can

be as large as 0.4 in group environments with $M_{\text{host}} = 10^{13} M_{\odot}$. F_{strip} in small and compact groups with low IGM temperature ($T_{\text{IGM}} = 10^6$ K) can be as large as ~ 0.6 , if the groups have IGM with $\rho_{\text{IGM}} \approx 10^5 M_{\odot} \text{ kpc}^{-3}$ and $V_r \approx 400 \text{ km s}^{-1}$. These results imply that strangulation can happen even in small and/or compact groups with higher velocity dispersions.

(3) The presence of disk gas in disk galaxies can suppress the ram pressure stripping of their halo gas owing to hydrodynamical interaction between halo and disk gas. This result suggests that stripping processes of halo gas by ram pressure can be significantly different between late- and early-type galaxies with and without disk gas.

(4) Disk galaxies with lower masses (M_{gal}) are likely to show appreciably higher F_{strip} in a given environment. This result implies that strangulation can proceed more rapidly and more efficiently in disk galaxies with lower M_{gal} . The remaining halo gas of disk galaxies after ram pressure stripping shows more compact and flattened distributions and has jellyfish-like configurations, irrespective of M_{gal} .

(5) Most halo gas stripped by ram pressure in disk galaxies can originate from the outer parts of the halos so that only halo gas initially in the inner parts can be accreted onto the disks during/after ram pressure stripping. This suggests that strangulation is more efficient in outer disks and thus that truncation of star formation by stripping of halo gas may well proceed outside-in in disk galaxies. This outside-in truncation can provide a clue to the origin of red passive spirals in distant groups and clusters.

(6) Ram pressure stripping of galactic halo gas is much more efficient than that of disk gas in groups and clusters and efficiencies of halo and disk gas stripping can be different in different environments with different M_{host} , V_r , and ρ_{IGM} . Thus the mass ratios of halo gas to disk one in disk galaxies can reflect in what environments they have resided in their histories and thus be significantly different between disk galaxies.

(7) It is suggested that recycling processes of ISM and chemical evolution of galaxies significantly change after removal of galactic halo gas. One example is that gaseous ejecta from energetic massive OB stars and supernovae can more easily escape from disk galaxies after halo gas stripping owing to much less effective hydrodynamical interaction between gaseous ejecta and halo gas. Abundance patterns of ISM in disk galaxies are thus suggested to be significantly different between different environments.

(8) Small fractions of halo gas (0.3–1%) in disk galaxies can be accumulated into the central 1 kpc of galactic bulges owing to the compression of halo gas during ram pressure stripping. These gas can be used for fueling the central MBH and activating weak AGN there, though the final dynamical fate of the accumulated halo gas is beyond the scope of this paper.

6 ACKNOWLEDGMENT

I am grateful to the anonymous referee for valuable comments, which contribute to improve the present paper. KB acknowledges the financial support of the Australian Research Council throughout the course of this work. I am grateful to participants in the last conference “galaxy evo-

lution and environment” held in Kuala Lumpur on 30 March – 3 April 2009 for their having extensive discussion on galaxy evolution in groups and clusters with me, which helped me better interpret the present numerical results in terms of the corresponding observational ones. Numerical computations reported here were carried out on the GRAPE system at the University of New South Wales and that kindly made available by the Center for computational astrophysics (CfCA) of the National Astronomical Observatory of Japan.

REFERENCES

- Abadi, M. G., Moore, B., Bower, R. G., 1999, MNRAS, 308, 947
- Balogh, M. L., Morris, S. L., Yee, H. K. C., Carlberg, R. G., Ellingson, E., 1999, ApJ, 527, 54
- Balogh, M. L., Navarro, J. F., Morris, S. L., 2000, ApJ, 540, 113
- Balogh, M. L., et al., 2004, MNRAS, 348, 1355
- Bamford, S. P. et al., 2009, MNRAS, 393, 1324
- Bekki, K., 1998, ApJL, 502, 133
- Bekki, K., Cocuh, W. J., 2003, ApJL, 596, 13
- Bekki, K., Chiba, M., 2006, ApJL, 637, L97
- Bekki, K., Peng, E. W., 2006, MNRAS, 370, 1737
- Bekki, K., Couch, W. J., Shioya, Y., 2001, PASJ, 53, 395
- Bekki, K., Couch, W. J., Shioya, Y., 2002, ApJ, 577, 651
- Bekki, K., Tsujimoto, T., Chiba, M., 2009, ApJ, 692, L24
- Binney, J., Tremaine, S. 1987 in Galactic Dynamics, Princeton; Princeton Univ. Press.
- Couch, W. J., Barger, A. J., Smail, I., Ellis, R. S., Sharples, R. M., 1998, ApJ, 497, 188
- Dressler, A., Smail, I., Poggianti, B. M., Butcher, H., Couch, W. J., Barger, A. J., Ellis, R. S., Oemler, A. Jr., 1999, ApJS, 122, 51
- Dressler, A., Rigby, J., Oemler, A., Fritz, J., Poggianti, B. M., Rieke, G., Bai, L., 2009, ApJ, 693, 140
- Font, A. S. et al., 2008, MNRAS, 389, 1619
- Freeman, K. C., 1970, ApJ, 160, 811
- Hester, J. A., 2006, ApJ, 647, 910
- Jeltema, T. E., Binder, B., Mulchaey, J. S., 2008, ApJ, 679, 1162
- Kapferer, W., Sluka, C., Schindler, S., Ferrarri, C., Ziegler, B., 2009, A&A, 499, 87
- Kawata, D., Mulchaey, J. S., 2008, ApJL, 672, 103
- Kronberger, T., Kapferer, W., Ferrari, C., Unterguggenberger, S., Schindler, S., 2008, A&A, 481, 337
- Larson, R. B., Tinsley, B. M., Caldwell, C. N., 1980, ApJ, 237, 692
- Marcolini, A., Brighenti, F., D’Ercole, A., 2003, MNRAS, 345, 1329
- McCarthy, I. G., et al., 2008, MNRAS, 383, 593
- Navarro, J. F., Frenk, C. S., White, S. D. M., 1996, ApJ, 462, 563 (NFW)
- Padmanabhan, T., 1993 Structure Formation in the Universe, Cambridge University Press
- Poggianti, B. M., Smail, I., Dressler, A., Couch, W. J., Barger, A. J., Butcher, H., Ellis, R. S., Oemler, A. Jr., 1999, ApJ, 518, 576
- Poggianti, B. M., et al., 2008, ApJ, 684, 888




Figure 1. Morphological evolution of disk gas projected onto the y - z plan in a disk galaxy for the standard model. The time T in units of Gyr is shown in the upper left corner for each frame.

Rasmussen, J., Ponman, T. J., Verdes-Montenegro, L., Yun, M. S. Borthakur, S., 2008, MNRAS, 388, 1245
 Roediger, E., Brüggén, M., 2008, MNRAS, 388, L89
 Sellwood, J. A., Carlberg, R. G., 1984, ApJ, 282, 61
 Sembach, K. R. et al., 2003, ApJS, 146, 165
 Sugimoto, D., Chikada, Y., Makino, J., Ito, T., Ebisuzaki, T., Umemura, M., 1990, Nat, 345, 33
 Tonnesen, S., Bryan, G. L., 2008, ApJL, 684, 9
 Vollmer, B., Soida, M., Otmianowska-Mazur, K., Kenney, J. D. P., van Gorkom, J. H., Beck, R., 2006, A&A, 453, 883
 Wielen R., 1977, A&A, 60, 263
 Wilman, D. J., Oemler, A., Mulchaey, J. S., McGee, S. L., Balogh, M. L., Bower, R. G., 2009, ApJ, 692, 298




Figure 2. The number distributions of gaseous pressure (P) for disk gas particles at $T = 0.21$ Gyr (blue) and $T = 0.56$ Gyr (red) in the standard model.

APPENDIX A: PHYSICAL PROPERTIES OF GAS DISKS DURING RAM PRESSURE STRIPPING

Fig. A1 shows the time evolution of morphological properties of the gas disk for the last 0.56 Gyr in the standard model. As the halo gas is compressed by the strong ram pressure of IGM, hydrodynamical interaction between disk and halo gas becomes stronger ($T = 0.21$ Gyr). Gaseous spiral arms can be developed in the inner region of the disk during ram pressure stripping, though the outer part of the gas disk can lose a minor fraction of its initial gas ($T = 0.28$ Gyr). The final morphology of the gas disk appears to be more compact ($T = 0.56$ Gyr) owing to compression of the gas disk by the halo gas and IGM.

Fig. A2 shows the number distributions of pressure of disk gas particles (P) at $T = 0.21$ Gyr and $T = 0.56$ Gyr in the standard model. The mean P at $T = 0.21$ Gyr and $T = 0.56$ Gyr are 1.1×10^{-11} dyn cm $^{-2}$ and 1.3×10^{-11} dyn cm $^{-2}$, respectively. (i.e., mean $\log_{10} P$ at $T = 0.21$ Gyr and $T = 0.56$ Gyr are -10.94 and -10.88 , respectively). Although the mean pressure is not so different between the two time steps, the mass fraction of disk gas particles that have pressure higher than 5×10^{-11} dyn cm $^{-2}$ (f_{thres}) is significantly different: $f_{\text{thres}}=0.04$ at $T = 0.21$ Gyr and 0.09 at $T = 0.56$ Gyr. This would reflect the fact that as time passes by, the disk gas particles can feel stronger pressure of halo gas (and IGM) owing to the more strongly compressed halo gas.

Recently Kapferer et al. (2009) have shown that if the pressure of IGM is as high as 5×10^{-11} dyn cm $^{-2}$, ram pressure can enhance significantly star formation in disk galaxies owing to compression of disk gas by ram pressure force. Our results in Fig. A2 therefore imply that star formation can be enhanced for some minor fraction ($< 10\%$) of disk gas during ram pressure stripping. Although numerical models P in Kapferer et al. (2009) and in the preset study are different, our results in Fig. A2 confirm the early suggestion (e.g., Bekki & Couch 2003; Kronberger et al. 2008) that ram pressure can trigger formation of new stars in gas disks. Numerical results by Kronberger et al. (2008) and by the present study are consistent with each other in that they show more efficient stripping of disk gas in outer parts of gas disks: star formation might well be truncated outside-in.

APPENDIX B: DESCRIPTION OF THE ANIMATION

We have made a simple animation for the time evolution of mass distributions of halo and disk gas projected onto the x - z plane for the standard model. Only halo and gas particles are shown by magenta and cyan, respectively for clarity: Dark matter halo, stellar disk and bulge, and hot IGM are not shown. The model parameters for the standard model are described in detail in the main text.

Hot gas particles for IGM are initially placed on equally spaced grids so that the adopted uniform density distribution of IGM can be achieved in the present study. In the early phase of ram pressure stripping of galactic halo gas by IGM, some grid-like local structures can be seen owing to (i) the adopted IGM distribution and (ii) smaller number of the IGM particle. Since these local structures can soon disappear, the final mass and distributions of galactic halo gas after ram pressure stripping, which are the main subjects of the present paper, can not be influenced significantly by the development of such structures (which is due to the adopted numerical models for IGM). The movie is available with the online version of the paper.